

---



---

Neurology

---



---

# The Post-Polio Syndrome: Current Concepts and Treatment

Sidney A. Spector, MD, PhD,\* Kumaraswamy Sivakumar, MD†

\*Neurological Physicians of Arizona, Inc., Mesa, Ariz.

†National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, Md.

The post-polio syndrome (PPS) occurs in clinically stable individuals decades after recovery from an initial attack of paralytic poliomyelitis; it is characterized by new muscular weakness and atrophy and generalized fatigue. PPS is believed to occur when muscles originally affected by paralytic polio begin to lose their innervation and weaken as a result of the death of remaining anterior horn cells of the spinal cord. Although it has been suggested that immune dysregulation or persistence of chronic polio infection plays a role, it is more likely that motor neuron attrition is the result of metabolic dysfunction. Neurotrophic factors that enhance the stability and growth of motor neurons and their synaptic connections are being studied as a means to stall or reverse the progression of weakness and atrophy that occurs in PPS.  
[Infect Med 14(6):462,465,466,471,472,474-478, 1997]

Key words: Neuromuscular diseases • Enterovirus infections • Poliomyelitis, paralytic • Post-poliomyelitis syndrome

**T**he World Health Organization has set the year 2000 as the target date for the global eradication of polio.<sup>1</sup> Although use of the live vaccine in the Western Hemisphere continues to cause a handful of cases each year, the risk of contracting polio remains mainly through importation of the disease from areas of the world where it is

endemic. Even in those areas, dramatic declines have occurred over the past 5 years. Yet the debilitation associated with polio is felt anew by some clinically stable individuals decades after their initial episode of paralytic poliomyelitis, with the onset of progressive muscular weakness known as the post-polio syndrome (PPS).

*Dr. Spector is Associate Director of the Neuromuscular Disease Section at Neurological Physicians of Arizona, Inc., Mesa, Ariz. Dr. Sivakumar is Staff Research Associate in the Neuromuscular Diseases Section, Medical Neurology Branch, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, Md.*

## *Epidemiology of Polio*

The path to the successful eradication of poliomyelitis was forged by the introduction of the inactivated polio vaccine of Salk in 1955 and the live attenuated oral polio vaccine of Sabin in 1961.<sup>2</sup> Since 1979, no case of paralytic poliomyelitis caused by the wild-type virus has been reported in the Western Hemisphere.<sup>3</sup> Throughout the world, however, poliomyelitis continues to occur due to live poliovirus and other enterovirus pathogens (coxsackie-

---

See editorial on page 438

---

virus types A and B, echoviruses, and enterovirus types 30 and 71),<sup>4</sup> live vaccine,<sup>5</sup> infection of nonimmunized groups,<sup>6</sup> infection of inadequately vaccinated populations,<sup>7</sup> and intramuscular injections temporally related to poliovirus infection or administration of the live polio vaccine.<sup>8</sup>

## *Development of PPS*

Shortly after the development of the acute febrile illness, a small percentage of individuals infected by the poliovirus will develop varying degrees of limb and bulbar

*continued on page 465*

continued from page 462

analysis according to the extent of injury and death of motor neurons. The motor neurons that survive the initial attack of the poliovirus go on to reinnervate originally denervated muscle fibers during a period of recovery typically lasting months to years,<sup>9</sup> and they maintain effective neuromuscular connections with functional stability for many years.<sup>9</sup> Then, decades later, some of these paralytic polio patients develop symptoms of PPS.

Although cases of patients with PPS were initially described more than 100 years ago, only recently have these new symptoms and signs been directly attributed to the paralytic disease.<sup>10-13</sup> While infection with the poliovirus leads to paralytic poliomyelitis in only 0.1% to 1% of patients,<sup>14</sup> more than half of individuals who do develop paralytic poliomyelitis have been reported to have the new muscular weakness characteristic of PPS.<sup>15-17</sup>

#### Clinical Characteristics of PPS

PPS comprises a constellation of symptoms, including new muscle weakness, atrophy, and fatigue; generalized fatigue; and muscle and joint pain. These symptoms develop 10 to 40 years after the initial attack of paralytic poliomyelitis,<sup>13,16-22</sup> (Table I), with the greatest incidence occurring 30 to 40 years after the initial infection.

Because virologic confirmation of polio infection was not available until the 1950s, older individuals describing these symptoms should provide a history of a well-documented, acute febrile illness during childhood or adolescence; preferably, this febrile illness should be associated temporally with a polio epidemic and should have resulted in partial or complete paralysis of a limb or bulbar musculature (ie, of the lips, tongue, pharynx, and larynx), followed by some degree of recovery of motor function, with

functional stability lasting for at least 10 years.<sup>23</sup> The clinical presentation of residual muscle atrophy, weakness, and areflexia in at least 1 limb with normal sensation, accompanied by histopathologic and electromyographic findings of chronic denervation, support the diagnosis of previous paralytic poliomyelitis. Within this clinical setting, presentation of newly progressive weakness, atrophy, muscle or joint pain, and fatigue suggests the diagnosis of PPS. However, the clinician must exclude other known medical, neurologic, orthopedic, and psychiatric disorders that may help to explain the development of the patient's new symptoms, such as back injuries, radiculopathies, compression neuropathies, degenerative arthritis, hyper- or hypothyroidism, fibromyalgia, anxiety, and depression.

#### Post-Polio Muscular Atrophy

Of all of the individuals diagnosed with poliomyelitis, only a small percentage of patients actually develop new symptoms of slowly progressive muscle weakness and atrophy, with or without fatigue, after many years of functional stability.<sup>15-18,22,24,25</sup> These symptoms are attributable to post-polio progressive muscular atrophy (PPMA), the further deterioration of remaining motor neurons and their collateral neuromuscular connections. The new weakness is usually randomly distributed and asymmetric, and it may involve portions of 1 or more muscles of any limb. Although it appears that any muscle is ultimately vulnerable, including those without previous clinical involvement, muscles previously affected by the original attack of polio, whether fully or partially recovered, are more likely to become involved than muscles that were clinically unaffected by the original infection.<sup>26</sup> As the remaining previously stable muscles be-

**Table I**  
Clinical Features of  
Post-Polio Syndrome

#### Neuromuscular

PPMA (new weakness of appendicular, bulbar, or respiratory muscles; new muscular atrophy; myalgias; fasciculations)

#### Muscle fatigue

New respiratory difficulties

#### Sleep apnea

#### Musculoskeletal

Joint pain

Joint and skeletal deformities

Decreased endurance

#### General

Generalized fatigue

Intolerance to cold

Increased sleep requirements

Psychologic problems

PPMA = post-polio muscular atrophy.

come progressively weaker, daily activities—which, for many years after recovery from acute paralytic poliomyelitis, may have been accomplished by remarkable compensatory improvisation—become increasingly more difficult to perform. In some of these patients, intrinsic muscle pain, cramps, and coarse fasciculations—indications of an ongoing neurogenic process—may either precede or be associated with the new muscle weakness.<sup>19,25,27</sup>

**Dysphagia.** In addition to involvement of appendicular muscles, new weakness of bulbar muscles may develop in patients with PPMA.<sup>28,29</sup> Although only 10% to 20% of polio patients actually complain of residual dysphagia, it appears that nearly all individuals with PPMA evaluated with videofluoroscopy demonstrate asymmetric weakness of pharyngeal or laryngeal muscles; impaired tongue movements; pooling in the valleculae or pyriform sinuses; or, rarely, aspiration.<sup>30</sup> Patients commonly complain of food or pills sticking in the throat, and intermittent choking on food. When present, symptoms of dysphagia are progressive and more severe in patients who have had

PPS *continued*

previous clinical bulbar involvement, although subclinical dysfunction may be detected in bulbar muscles clinically spared by the initial illness.<sup>29</sup>

**Respiratory symptoms.** Progressive respiratory insufficiency is seen in patients left with diminished respiratory reserve after the acute attack of polio.<sup>31,32</sup> In most cases, patients functioning with marginal respiratory capacity (eg, those who become easily tired or short of breath on exertion), who may have been weaned successfully from ventilator use after recovery from the acute illness, ultimately develop progressive respiratory difficulties.<sup>32</sup> Respiratory insufficiency directly related to PPMA is usually caused by residual diaphragmatic muscle weakness, or, more rarely, occurs as a result of central hypoventilation from bulbar involvement of the reticular activating system (a network of neurons in the spinal cord, brain stem, and thalamus that controls many unconscious motor activities) and sleep regulatory centers. Secondary causes also may contribute, including progressive scoliosis, pulmonary emphysema, cardiovascular insufficiency, and inactivity.<sup>33</sup>

Sleep apnea is often found in PPMA patients in association with other symptoms of respiratory compromise and residual bulbar dysfunction.<sup>32,34,35</sup> The central component of this common problem is dysfunction of the reticular formation. Peripheral factors include progressive weakness of diaphragmatic, intercostal, abdominal, and pharyngeal muscles; obesity; emphysema; and musculoskeletal deformities.

**Fatigue.** The majority of patients with PPS, and nearly all patients with PPMA, experience fatigue and diminished stamina. Patients may describe a generalized, overpowering feeling of tiredness and malaise unrelated to physical ac-

tivity, or may feel a more specific sensation of fatigue localized to muscles. In contrast to patients with multiple sclerosis (MS) or chronic fatigue syndrome (CFS), who frequently experience a generalized malaise throughout the day, PPS patients usually develop mid-day fatigue that is ameliorated by brief rest periods. Psychopathologic symptoms of anxiety, depression, and stress are not cofactors in the development of fatigue in patients with PPS.<sup>36</sup>

Many survivors of paralytic poliomyelitis develop new musculoskeletal symptoms,<sup>15,22,37-40</sup> including diminished functional aerobic capacity, decreased endurance, and weight gain; these symptoms are combined with degenerative arthri-

*Childhood surgical intervention—including tendon transfers, joint functions, and shortening of the long bones of the lower extremities—may unpredictably alter the natural biomechanics of joints during subsequent growth and recovery from paralysis.*

tis, scoliosis, and pain in joints that have been biomechanically disadvantaged or deformed owing to long-standing asymmetry of size and strength of previously affected muscles. Although these new musculoskeletal deficits are often a prominent feature of PPS, there is no evidence that they represent neuromuscular sequelae of prior polio. Rather, these new symptoms are invariably the result of secondary processes such as long-standing scoliosis, obesity, and childhood surgical interventions. The latter—including tendon transfers, joint fusions, and shortening of the long bones of the lower extremities<sup>40</sup> intended to improve mobility and stability after recovery from the acute

paralytic illness—may unpredictably alter the natural biomechanics of joints during subsequent growth and recovery from paralysis, resulting in excessive wear and tear on joints, ligaments, and tendons. New musculoskeletal complaints in patients with prior paralytic poliomyelitis, directly or indirectly attributable to prior polio, should be considered as elements of PPS, although they are not neuromuscular in origin (Table I).

#### *Epidemiology of PPS*

There are more than 1.63 million survivors of paralytic poliomyelitis.<sup>23</sup> Between 20% and 60% of these individuals may develop PPS,<sup>15,23,41</sup> but when individuals with musculoskeletal symptoms are excluded, the percentage of paralytic poliomyelitis patients who develop PPMA appears to be much lower.<sup>10,42</sup> Individuals more likely to develop PPS include older patients<sup>15,43</sup> who have tended to develop more severe weakness during the acute paralytic illness<sup>34,41</sup> and have experienced greater residual deficits,<sup>26</sup> as well as those presenting after longer periods of stability.<sup>43</sup> The interval of clinical stability ranges from 8 to 71 years,<sup>18</sup> with the highest incidence of PPS occurring 30 to 40 years after the acute polio attack.<sup>18,19,34</sup>

The natural course of the symptoms of fatigue and pain of PPS is difficult to assess because these complaints are not amenable to objective assessment. On the other hand, the power of muscles weakened by PPMA may be quantified, and it appears to diminish at a variable but slow rate of about 1% per year,<sup>25</sup> although the patient may experience periods of relative stability lasting many years.<sup>42,44,45</sup> Slowly progressive loss of strength of the tongue and swallowing muscles appears to occur more consistently.<sup>24</sup>

*continued on page 471*

continued from page 466

**Pathogenesis of PPMA.** The cause of new anterior horn-cell degeneration and denervation in muscles of patients with PPMA is unknown. The development of PPMA in younger patients cannot be explained by aging, which normally leads to gradual motor-neuronal loss only in individuals above the age of 60.<sup>46</sup> More likely, after many years of stability, the chronically increased metabolic burden placed on the neuron becomes too great to sustain all of its synapses, leading to a slow attrition of sprouts and the ultimate death of the motor neuron.<sup>18,47</sup> However, recent studies have raised the possibility that immune dysregulation or persistence of chronic polio infection may be responsible for new neuro-muscular symptoms.

During the initial paralytic attack by the poliovirus, muscles are weakened to varying degrees, depending on the extent of injury or death of anterior horn cells of the spinal cord.<sup>48,49</sup> Whether muscles recover partially or completely depends on the ability of injured, healing motor neurons and normal, unaffected neurons to generate distal axonal sprouts to neighboring muscle fibers that have lost their innervation.<sup>50-52</sup>

**Neuronal findings.** Electrophysiologic recordings demonstrate abnormally large motor units of prolonged duration, both in weakened and clinically uninvolved muscles of polio patients, as remaining motor neurons reinnervate up to 5 times their normal complement of muscle fibers.<sup>26,50-54</sup> During a prolonged period of relative stability, new axonal sprouting occurs in previously uninvolved or recovered neurons as other motor neurons that can no longer sustain the increased numbers of connections begin to die.<sup>32</sup> Electrical studies of individual fibers innervated by a single motor axon in muscle of pa-




Figure 1. Remodelling of motor units during acute polio and early and late post-polio muscular atrophy (PPMA). Under normal conditions, anterior horn cells innervate set complement of muscle fibers in particular muscle. Acute paralytic poliomyelitis leads to injury or death of some of these cells. Remodelling of remaining motor neurons occurs during stable post-polio with effective reinnervation of newly denervated muscle fibers. Greater numbers of muscle fibers become denervated with progression of PPMA, resulting in presence of denervated atrophic muscle fibers in groups. Adapted from *Advances in Neurology: Amyotrophic Lateral Sclerosis* (1991; p 506). Copyright © 1991. Lippincott-Raven Healthcare.<sup>33</sup>

tients with either stable polio or PPMA demonstrate a high variability of impulse conduction (jitter), which continues indefinitely.<sup>51-53,55</sup> Jitter is not seen in normally innervated motor units and suggests the presence of ongoing denervation. Nerve cell adhesion molecules (N-CAMs), expressed on the surface of muscle fibers only after they lose innervation, are present in both stable and weakening post-poliomyelitis muscles,<sup>52,56</sup> providing a histologic correlate of remodeling of the motor unit. New muscle weakness and atrophy occur when the attrition of these neurons outweighs the capacity of the remaining anterior horn cells to sprout and innervate newly denervated muscle fibers, which is reflected histologically by increasing numbers of N-CAM-positive, denervated muscle fibers (Fig. 1).

**Immunologic findings.** Immunologic abnormalities have been observed in a fraction of patients with PPMA, including mild inflammation consisting of CD8+ and CD4+ cells surrounding healthy muscle fi-

bers that express MHC class I antigens on their surface; this inflammation is similar to that seen in immune-mediated myopathies,<sup>57</sup> inflammation in the parenchyma of gray matter of spinal cord,<sup>58,59</sup> intrathecal oligoclonal IgG bands,<sup>24</sup> and activated T cells in peripheral blood.<sup>60</sup> It has been suggested that immunologic dysregulation may play a role in the development of new symptoms of weakness and atrophy due to motor neuron degeneration,<sup>24,34,56</sup> although direct evidence for this theory is not yet forthcoming.

Although poliovirus is lytic and produces a monophasic illness,<sup>4</sup> an intriguing proposed hypothesis postulates that defective poliovirus particles or mutant viruses, having escaped detection by the immune system by persisting indefinitely in motor neurons that did not lyse, are subsequently reactivated to cause the death of these cells.<sup>56,61,62</sup> This hypothesis has been supported by the presence of intrathecal poliovirus-specific IgM and IgG antibodies, and high IgM antibody titers to

**Table II**  
**Differential Diagnoses**

- Degenerative arthritis:** Patients develop joint pain, without muscle atrophy or fasciculations; serum muscle enzymes are normal; electrodiagnostic and muscle biopsy studies do not demonstrate features of denervation.
- Fibromyalgia:** Patients experience generalized muscle soreness, fatigue, and malaise in the absence of clinical evidence of muscle atrophy or fasciculations, elevation in serum muscle enzymes, and normal electrodiagnostic and muscle biopsy evaluations.
- Radiculopathy:** Patients experience new signs of asymmetric muscle weakness, atrophy, and fasciculations. Associated symptoms of pain and paresthesias are not typically experienced by post-polio syndrome (PPS) patients. Neuroimaging of cervical or lumbar spine excludes degenerative disc disease or spinal stenosis.
- Multiple sclerosis (MS) or chronic fatigue syndrome (CFS) as causes of fatigue:** MS or CFS patients frequently experience malaise throughout the day; PPS patients usually develop midday fatigue ameliorated by brief periods of rest. MS is accurately diagnosed by typical central nervous system MRI and cerebrospinal fluid studies. In contrast to patients with PPS, patients with electrodiagnostic evaluations are typically normal in patients with CFS. Psychopathologic symptoms of anxiety, depression, and stress are not cofactors in the development of fatigue in patients with PPS, as they may be in patients with MS and CFS.
- Compression peripheral neuropathies:** Impingement of peripheral nerves of upper or lower extremities, resulting from long-standing musculoskeletal deformities or other unrelated processes, leads to asymmetric muscle atrophy. Pain and paresthesias are usually associated symptoms. Electrodiagnostic testing differentiates these peripheral neurogenic processes from the motor neuron disjunction of PPS.
- Acquired primary myopathy (eg, inclusion-body myositis [IBM]):** Like PPS, IBM advances slowly and asymmetrically, and there may be electromyographic abnormalities resulting from secondary distal axonal degeneration. PPS is excluded by muscle biopsy demonstrating typical inflammatory and structural histopathologic features of IBM (eg, rimmed vacuoles, endomysial inflammation, variable fiber size, and eosinophilic cytoplasmic inclusions).
- Amyotrophic lateral sclerosis (ALS):** ALS begins focally but progresses diffusely to involve bulbar and respiratory muscles, nearly always producing hyperreflexia, spasticity, and plantar extensor responses. These upper motor neuronal signs are only rarely discovered in PPS.
- Progressive muscular atrophy (a variant of ALS):** Consider whether there is a lack of upper motor neuronal involvement. Distinguished from PPS by rapid progression of atrophy. Occasionally, distinction may be made by muscle biopsy, which may reveal abnormally large fiber-type grouping in PPS.
- Multifocal motor neuropathy disease of peripheral nerves:** Slowly progressive asymmetric weakness and atrophy develop without changes in sensation. PPS is excluded by presence of characteristic electrodiagnostic nerve conduction abnormalities and demonstration of high titers of antiganglioside antibodies in the blood.

nosis of PPS is not difficult in an individual with a previously documented history of paralytic poliomyelitis who subsequently develops new musculoskeletal symptoms, signs, and weakness, and in whom other neurologic, orthopedic, and medical disorders have been excluded—these include radiculopathies, compression neuropathies, myopathy, fibromyalgia, and arthritis (Table II). For patients in whom the diagnosis of paralytic poliomyelitis is questionable, electrodiagnostic and histopathologic indications provide evidence of previous paralytic poliomyelitis.

On occasion, however, other neuromuscular processes must be considered when a well-documented history of polio is not forthcoming or when the pace of neuromuscular deterioration is rapid. For example, amyotrophic lateral sclerosis (ALS), a relatively rapidly progressing fatal disease of the motor neurons,

may begin focally, but it progresses diffusely to involve bulbar and respiratory muscles and nearly always produces hyperreflexia, spasticity, and plantar extensor responses—upper motor neuronal signs that rarely are found in PPS.<sup>18</sup> In the absence of upper motor neuronal involvement, the clinician should consider progressive muscular atrophy, a variant of ALS that may be distinguished from PPS only by the rapid progression of that disease. PPS may sometimes be confused with multifocal motor neuropathy,<sup>73,74</sup> a disease of the peripheral nerves in which slowly progressive asymmetric weakness and atrophy develop without changes in sensation. However, the presence of characteristic electrodiagnostic nerve conduction abnormalities<sup>73,74</sup> and the demonstration of high titers of antiganglioside antibodies in the blood<sup>75,76</sup> of these patients exclude PPS.

On rare occasions, an individual diagnosed initially as having PPS actually may have been weakened by a long-standing acquired primary myopathy such as inclusion-body myositis,<sup>57,77</sup> which also advances slowly in an asymmetric manner and which may display neurogenic electromyographic features because of secondary distal axonal degeneration.<sup>78,79</sup> A muscle biopsy demonstrating typical inflammatory and structural histopathologic features specific for inclusion-body myositis (eg, rimmed vacuoles, endomysial inflammation, variable fiber size, and eosinophilic cytoplasmic inclusions) will confirm the diagnosis of this myopathy.<sup>77</sup>

#### *Treatment*

Management of PPS addresses several aspects of the syndrome: muscular weakness, general fatigue, cardiovascular and respiratory

PPS *continued*

poliovirus in serum of a small number of patients with PPMA, suggesting a new immune response against poliovirus.<sup>63-65</sup> Further, amplification of poliovirus RNA particles from spinal fluid by polymerase chain reaction has revealed incomplete polioviral and other enteroviral sequences.<sup>65-67</sup> Although defective poliovirus particles that may be harbored in remaining anterior horn cells during many years of stability appear to become antigenic when ultimately released as these cells degenerate, there is no direct evidence for reactivation of virulent poliovirus in these patients.

**Pathogenesis of fatigue.** The cause of fatigue in patients with PPS appears to be multifactorial, involving both central and peripheral nervous system mechanisms. Reduced cortical drive to lower motor neurons<sup>68</sup> may play a role in the development of early central fatigue in patients with PPMA. In the periphery, neuromuscular fatigue has been attributed to retraction of axonal sprouts resulting from dying anterior horn cells, as suggested by the presence of increased jitter during single-muscle-fiber electrical recordings, histologic changes of neuromuscular synapses of affected muscles, and the transient amelioration of fatigue in PPMA patients treated with anticholinesterase inhibitors.<sup>69,70</sup> However, no neuromuscular transmission defect in PPMA patients was confirmed in assessment of functional performance and energy utilization of muscles during low-intensity aerobic exercise; rather, it appears that neuromuscular fatigue lies at a point beyond muscle fiber membrane excitation.<sup>71</sup> Histologic abnormalities of oxidative energy metabolism have been shown in clinically affected or asymptomatic muscles of PPMA patients. In addition, during graded aerobic exercise, patients with PPMA showed earlier development of fatigue in

clinically unaffected muscles, resulting from impaired oxidative metabolism; patients required a greater dependence on glycolytic energy pathways, as earlier development of intracellular metabolic acidosis rapidly depleted their energy stores of phosphocreatine.<sup>72</sup>

#### *Evaluation of Patients*

There are no laboratory parameters that specifically confirm the presence of PPS, and as such, this diagnosis is based on clinical grounds after other medical, orthopedic, neurologic, and psychiatric diseases have been excluded (Table I). Evidence of progression of weakness may be derived from annual evaluations<sup>25</sup> of strength of

*There are no laboratory parameters that specifically confirm the presence of PPS.*

affected muscle groups. Quantitative isometric or dynamic studies of muscle strength are superior to Medical Research Council manual muscle testing in assessing changes in strength over time. Swallowing studies may provide evidence of progressive bulbar impairment. Serial evaluations of photographs may reveal changes in muscle girth, while muscle size may be studied sequentially with computer tomographic or magnetic resonance imaging techniques.

Routine blood tests are usually normal, although muscle creatine kinase may be elevated to 1000 IU/L in up to 10% of patients with PPMA.<sup>24,66</sup> Elevation in cerebrospinal fluid protein without pleocytosis may be observed occasionally.<sup>18</sup> A small percentage of individuals with PPMA demonstrate high IgM antibody titers to poliovirus in serum,<sup>63</sup> as well as elevations of IgM oligoclonal bands specific to poliovirus intrathecally.<sup>64</sup>

Electrodiagnostic evaluations of muscle do not distinguish between residual paralytic poliomyelitis and new symptoms of weakness and atrophy of PPS, but these tests may confirm previous paralytic poliomyelitis in an otherwise unsubstantiated case.<sup>42,49-51</sup> Muscles that were either clinically affected or asymptomatic during the acute phase of polio, and those experiencing new symptoms, may display evidence of abnormal spontaneous activity; this activity appears in the form of fibrillations and positive waves, and abnormally large motor units indicative of remodelling of remaining motor units. Both of these features are indistinguishable from those in patients with previous polio or other neurogenic processes. Similarly, findings of single-motor-unit, single-muscle-fiber, and repetitive-stimulation studies cannot distinguish unequivocally between stable residual polio and PPMA. Nevertheless, these studies do confirm the presence of acute and chronic denervation, and they may be useful to exclude compression neuropathies involving peripheral nerves of the upper and lower extremities, which result from long-standing musculoskeletal deformities or other unrelated neurogenic processes.

While histologic evaluations are not diagnostic of PPS, they may provide evidence of prior polio and support the presence of ongoing neuromuscular remodelling. Neurogenic and myopathic changes are found to varying degrees in biopsied muscles that were previously weakened or unaffected during acute poliomyelitis, and are stable or are now developing new weakness. These changes include large fiber type grouping; groups of atrophic fibers; small, scattered angulated fibers; increased connective tissue; variation of fiber size; and occasional necrotic fibers.<sup>19,24</sup>

**Differential diagnosis.** The diag-

*continued on page 474*

blems, and joint pain. (See Table III for treatment summary, Table 1, "Resources for PPS Treatment and Support.")

**Muscular weakness.** Several studies indicate that muscle weakness in patients with PPMA may improve significantly with strengthening.<sup>80-82</sup> This remains a central issue in neurorehabilitation<sup>83</sup> because earlier uncontrolled studies suggested that excessive exercise in patients with reduced innervation could lead to further damage.<sup>84,85</sup> Nevertheless, it was demonstrated recently that patients with PPMA who completed a 10-week nonfatiguing, progressive-resistance, strength-training program significantly increased their strength in both symptomatic and minimally unaffected muscles—without serologic or histologic evidence of muscle damage.<sup>80</sup> Strength was maintained in nearly all muscles up to 5 months after cessation of training. The additional benefits of decreasing total fat mass<sup>86</sup> while enhancing bone mineral density<sup>87</sup> with strength training suggest that a program of supervised, nonfatiguing, progressive strength-training may be an efficacious therapeutic modality in patients with PPMA.

The swallowing difficulties encountered by patients with PPMA may be alleviated by postural adjustments, such as dropping the chin or turning the head to one side to minimize the deficit of a unilaterally weakened pharynx.<sup>30</sup> These modifications are best instituted by a speech therapist with special training in dysphagia.

**General fatigue.** The generalized fatigue experienced by most patients with PPS is alleviated to a great extent with frequent periods of rest, lasting between 15 minutes and 2 hours, after which they are able to resume their activities.<sup>26</sup> Clinical studies using prednisone,

**Table III**  
Treatment of the Post-Polio Syndrome

**Post-polio progressive muscular atrophy (PPMA)**

- Progressive-resistance, strength-training program for 10 weeks.
- For dysphagia: Speech therapists can help PPMA patients improve swallowing by teaching them postural adjustments, such as dropping the chin or turning the head to one side, to compensate for a unilaterally weakened pharynx.
- Pyridostigmine may be useful for neuromuscular fatigue.

**General fatigue**

- Frequent periods of rest lasting from 15 minutes to 2 hours.
- Treatment with prednisone, amantadine, insulin-like growth factor-1, or alpha-2 recombinant interferon has not resulted in consistent improvement of fatigue symptoms.

**Respiratory or cardiovascular problems**

- Aerobic exercise training can significantly enhance oxygen uptake and cardiovascular endurance. Walking on a treadmill, riding a stationary bicycle, and swimming or other pool exercises increase aerobic capacity.

**Muscle and joint pain and instability**

- Joint pain: analgesics, anti-inflammatory agents, and physical therapy. Gait training, changes in posture, weight reduction, and adjustments to orthotic devices (wheelchair, crutches, orthoses) also improve symptoms. Rarely, surgical repair of unstable joints may be required.
- Aerobic activity also reduces total body fat and improves movement efficiency and activities of daily living.

**Resources for PPS Treatment and Support**

Post-Polio Syndrome Research Center, NYU Rehabilitation Center  
New York University Medical Center  
400 E 34th Street, New York, NY 10016  
(212) 263-7300

Post-Polio Rehabilitation and Research Service  
Kessler Institute for Rehabilitation  
300 Market Street, Saddle Brook, NJ 07663-5309  
(201) 587-8500

International Polio Network  
5100 Oakland Avenue, #206, St. Louis, MO 63110  
(314) 534-0475

Polio Survivors Association  
12720 La Reina Avenue, Downey, CA 90242  
(310) 862-4508

National Center for Medical Rehabilitation Research  
National Institutes of Health  
Department of Health and Human Services  
Executive Plaza South, Room 450 West, 6120 Executive Boulevard  
Rockville, MD 20885  
(301) 402-2242

amantadine, recombinant insulin-like growth factor-1, or alpha-2 recombinant interferon have not demonstrated any consistent improvement in levels of fatigue in patients with PPS, although in in-

dividual cases some benefit may be obtained.<sup>88,89</sup> Preliminary study indicates that a component of peripheral neuromuscular fatigue in a small group of PPS patients may be ameliorated with anticholines-



## Related Articles on ...



Here are some of the most recent articles covering post-polio syndrome on Medscape ([www.medscape.com](http://www.medscape.com)):

- Lutwick LI, Chapnick EK: Bug of the Month: Weakness Upon Weakness
- CDC: Paralytic Poliomyelitis in the United States
- CDC: Poliomyelitis Prevention in the United States

You can find these and related articles by clicking on the "search" tab on the Medscape home page.

Medscape offers convenient, preconfigured key word searching of the thousands of full-text articles on the site. From within an article, simply click on a key word to initiate a search.

Free MEDLINE and AIDSLINE access is also available.



terases such as pyridostigmine.<sup>91</sup> A clinical trial studying these agents in a large population of PPS patients with fatigue is ongoing.

**Cardiovascular and respiratory problems.** Reduced cardiopulmonary function and capacity for oxygen by metabolically impaired muscles may lead to diminished endurance in patients with PPS.<sup>91,92</sup> All ambulatory patients with PPS should consider aerobic exercise training, which, in addition to significantly enhancing oxygen uptake and cardiovascular endurance,<sup>33,93</sup> also reduces total body fat and improves movement efficiency and activities of daily living.<sup>94</sup> Aerobic capacity can be increased by walking on a treadmill, riding a stationary bicycle, or swimming and participating in other pool exercises.

**Joint problems.** The management of the wide range of musculoskeletal symptoms of PPS patients has depended on supportive care and

symptomatic treatment. Joint pain is treated with analgesics, anti-inflammatory agents, and physical therapy. It is not uncommon to encounter patients with new joint pain whose symptoms are improved significantly with gait training, changes in posture, weight reduction, and adjustments to orthotic devices (wheelchair, crutches, orthoses).<sup>95</sup> Rarely, surgical reconstruction may be necessary to repair unstable joints.<sup>96</sup>

**New clinical trials.** Although no treatment is yet available to reverse the ongoing loss of anterior horn cells and their sprouts in patients with PPMA, clinical trials utilizing neurotrophic factors to stimulate nerve growth are underway. For example, in a study of developing rats, ciliary-derived neurotrophic factor was shown to attenuate the loss of motor neurons that occurs naturally during prenatal development<sup>97</sup> and also spares facial motor neurons in a mouse

model of progressive motor neuropathy.<sup>98</sup> Insulin-like nerve growth factor (IGF-1), a peptide synthesized in muscle, liver, and kidney in response to pituitary growth hormone, has been shown to stimulate anterior horn cells, in vitro and in experimental animals, to generate new axonal sprouts and synapses to muscle fibers.<sup>99</sup> Given that the primary problem in patients with PPMA is the progressive deterioration of remaining motor neurons, these proteins may have the clinical potential to maintain existing motor neurons and their connections. It is possible that such proteins could even stimulate the growth of new nerve terminals (sprouts) to form new synapses, in order to attenuate or reverse the progression of many of the debilitating neuromuscular symptoms of patients with PPMA.

### Summary and Conclusions

Of the estimated 1.63 million polio survivors in the US, half may develop PPS.<sup>23</sup> Onset of PPS and the characteristic muscular atrophy typically occurs 10 to 40 years after an initial attack of paralytic polio. Common symptoms include new muscular weakness, pain, and atrophy; dysphagia; respiratory insufficiency including sleep apnea; fatigue; and joint pain and arthritic changes. Exhaustion of remaining motor neurons and loss of innervation of previously asymptomatic muscles appear to be the primary cause of PPS, but immune dysregulation or persistence of chronic polio infection may be contributory. Although PPS is managed primarily by supportive and symptomatic care, nonfatiguing strengthening exercise may result in short-term improvements, and studies are under way to determine whether nerve-growth-stimulating factors may preserve existing motor neurons and their



connections to muscle. In addition to physical therapy, patients may benefit from psychological counseling as they learn to cope with this disabling and belated aftermath of their acute poliomyelitis. □

#### Drugs Mentioned in This Article

Amantadine	<i>Symmetrel</i>
Prednisone	Generic
Pyridostigmine	<i>Mestinon</i> , <i>Regonol</i>

#### References

- Lepow ML: Epidemiology of polio: US and worldwide. *Infect Med* 13(suppl D):S17-S25, 1996.
- Robbins RC: Eradication of polio in the Americas. *JAMA* 270:1857-1859, 1993.
- Certification of poliomyelitis eradication—the Americas. *MMWR Morb Mortal Wkly Rep* 43:720-722, 1994.
- Sabin AB: Poliomyelitis, in Horsfall FL Jr, Tamm J (eds): *Viral and Rickettsial Infections of Man*. Philadelphia, Lippincott, 1980, pp 1348-1381.
- Nkowane BM, Wassilak SF, Orenstein, WA, et al: Vaccine-associated paralytic poliomyelitis. *JAMA* 257:1335-1340, 1987.
- Centers for Disease Control and Prevention: CDC isolation of wild poliovirus type 3 among members of a religious community objecting to vaccination—Alberta, Canada, 1993. *MMWR Morb Mortal Wkly Rep* 42:337-339, 1993.
- Kim-Farley RJ, Rutherford G, Lichfield P, et al: Outbreak of paralytic poliomyelitis, Taiwan. *Lancet* 2:1322-1324, 1984.
- Strebel PM, Ion-Nedelcu N, Baughman AL, et al: Intramuscular injections within 30 days of immunization with oral poliovirus vaccine: A risk factor for vaccine-associated paralytic poliomyelitis. *N Engl J Med* 332:500-506, 1995.
- Mulder DW: Clinical observations on acute poliomyelitis, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 1-10.
- Mulder DW, Rosenbaum RA, Layton DO Jr: Late progression of poliomyelitis or forme fruste amyotrophic lateral sclerosis. *Mayo Clin Proc* 47:756-761, 1972.
- Kayser-Gatchalian MC: Late muscular atrophy after poliomyelitis. *Eur Neurol* 10:371-380, 1973.
- Alter M, Kurland LT, Molgaard CA: Late progressive muscular atrophy and antecedent poliomyelitis, in Rowland LP (ed): *Human Motor Neuron Diseases*. New York, Raven Press, 1982, pp 303-309.
- Halstead LS, Wiechers DO (eds): *Late Effects of Poliomyelitis*. Miami, Symposia Foundation, 1985.
- Morens DM, Pallansch MA, Moore M: Polioviruses and other enteroviruses, in Belshe R (ed): *Textbook of Human Virology*. St. Louis, Mosby Year-Book, 1991, pp 427-497.
- Codd MB, Mulder DW, Kurland LT, et al: Poliomyelitis in Rochester, Minnesota, 1935-1955: Epidemiology and long-term sequelae: A preliminary report, in Halstead LS, Wiechers DO (eds): *Late Effects of Poliomyelitis*. Miami, Symposia Foundation, 1985, pp 121-134.
- Halstead LS, Rossi DC: Postpolio syndrome: Clinical experience with 132 consecutive outpatients, in Halstead LS, Wiechers DO (eds): *Research and Clinical Aspects of the Late Effects of Poliomyelitis*. White Plains, NY, March of Dimes Birth Defects Foundation, 1987, pp 13-26.
- Cashman NR, Maselli R, Wollman RL, et al: New muscle atrophy as a late symptom of the post-poliomyelitis syndrome. *Clin Ecol* 5:11-13, 1987.
- Jubelt B, Cashman NR: Neurological manifestations of the post-polio syndrome. *Crit Rev Neurobiol* 3:199-220, 1987.
- Dalakas MC, Hallett M: The post-polio syndrome, in Plum F (ed): *Advances in Contemporary Neurology*. Philadelphia, FA Davis, 1988, pp 51-94.
- Munsat T (ed): *The Post Polio Syndrome*. Stoneham, UK, Butterworths, 1990.
- Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995.
- Dalakas MC: New neuromuscular symptoms in patients with old poliomyelitis: A three-year follow-up study. *Eur Neurol* 25:381-387, 1986.
- Jubelt B, Drucker J: Post-polio syndrome: An update. *Semin Neurol* 13:283-290, 1993.
- Dalakas MC: Morphological changes in the muscles of patients with post-poliomyelitis neuromuscular symptoms. *Neurology* 38:99-104, 1988.
- Dalakas MC, Elder G, Hallett M, et al: A long-term follow-up study of patients with post-poliomyelitis muscular symptoms. *N Engl J Med* 314:959-963, 1986.
- Dalakas MC: The post-polio syndrome as an evolved clinical entity, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 68-80.
- Fetell MR, Smallberg G, Lewis LD, et al: A benign motor neuron disorder: Delayed cramps and fasciculations after poliomyelitis or myelitis. *Ann Neurol* 11:423-427, 1982.
- Sonies BC, Dalakas MC: Dysphagia in patients with the post-polio syndrome. *N Engl J Med* 324:1162-1167, 1991.
- Buchholz DW, Jones B: Post-polio dysphagia: Alarm or caution? *Orthopedics* 14:1303-1305, 1991.
- Sonies BC, Dalakas MC: Progression of oral-motor and swallowing symptoms in the post-polio syndrome, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 87-95.
- Fischer DA: Poliomyelitis: Late respiratory complications and management. *Orthopedics* 8:891-894, 1985.
- Bach JR, Alba AS: Pulmonary dysfunction and sleep disordered breathing as post-polio sequelae: Evaluation and management. *Orthopedics* 14:1329-1337, 1991.
- Dalakas MC, Illa I: The post-polio syndrome: Concepts in pathogenesis and etiologies, in Rowland LP (ed): *Advances in Neurology: Amyotrophic Lateral Sclerosis*. New York, Raven Press, 1991, pp 495-511.
- Halstead LS, Wiechers DO, Rossi CD: Late effects of polio-myelitis: A national survey, in Halstead LS, Wiechers DO (eds): *Research and Clinical Aspects of the Late Effects of Poliomyelitis*. Miami, Symposia Foundation, 1985, pp 11-31.
- Guilleminault C, Motta J: Sleep apnea syndrome as a long-term sequela of poliomyelitis, in Guilleminault C (ed): *Sleep Apnea Syndromes*. New York, Kroc Foundation, 1978, pp 309-315.
- Clark K, Dinsmore S, Grafman J, et al: A personality profile of patients diagnosed with post-polio syndrome. *Neurology* 44:1809-1811, 1994.
- Agre JC, Rodriguez AA, Sperling KB: Symptoms and clinical impressions of patients seen in a postpolio clinic. *Arch Phys Med Rehabil* 70:367-370, 1989.
- Anderson AS, Levine SA, Gellert H: Loss of ambulatory ability in patients with old anterior poliomyelitis. *Lancet* 2:1061-1063, 1972.
- Cosgrove JL, Alexander MA, Kitts EL, et al: Late effects of poliomyelitis. *Arch Phys Med Rehabil* 68:4-7, 1987.
- Weinstein L: Diagnosis and treatment of poliomyelitis. *Med Clin North Am* (Sept):1377-1402, 1948.
- Windebank AJ, Litchy WJ, Daube JR, et al: Late effects of paralytic poliomyelitis in Olmsted County, Minnesota. *Neurology* 41:501-507, 1991.
- Windebank AJ, Litchy WJ, Daube JR, et al: Lack of progression of neurological deficit in survivors of paralytic polio: A 5-year prospective population-based study. *Neurology* 46:80-84, 1996.
- Trojan DA, Cashman NR, Shapiro S, et al: Predictive factors for post-poliomyelitis syndrome. *Arch Phys Med Rehabil* 75:770-777, 1994.
- Grimby G, Hedberg M, Henning G: Changes in muscle morphology, strength and enzymes in a 4-5 year follow-up of subjects with poliomyelitis sequelae. *Scand J Rehab Med* 26:121-130, 1994.
- Ivanyi B, Nelemans PJ, de Jongh R, et al: Muscle strength in postpolio patients: A prospective follow-up study. *Muscle Nerve* 19:738-742, 1996.
- Tomlinson BE, Irving D: The numbers of limb motor neurons in the human lumbosacral cord throughout life. *J Neurol Sci* 34:213-219, 1977.
- Dalakas MC: Pathogenetic mechanisms of post-polio syndrome: Morphological, electrophysiological, virological and immunological correlations, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 167-185.
- Bodian D: Poliomyelitis: Histopathologic basis of clinical findings. *Am J Med* 6:563-578, 1949.
- Bodian D: Poliomyelitis, in Minckler J (ed): *Pathology of the Nervous System*, New York, McGraw-Hill, 1984, pp 2323-2364.
- Bodian D: Motoneuron disease and recovery in experimental poliomyelitis, in Halstead LS, Wiechers DO (eds): *Late Effects of Poliomyelitis*. Miami, Symposia Foundation, 1985, pp 45-55.
- Wiechers DO, Hubbell SL: Late changes in the motor unit after acute poliomyelitis. *Muscle Nerve* 4:524-528, 1982.

- Cashman NR, Maselli RL, Wollmann R, et al: Late denervation in patients with antecedent paralytic poliomyelitis. *N Engl J Med* 317:7-12, 1987.
- Ravits J, Hallett M, Baker J, et al: Clinical and electromyographic studies of post-poliomyelitis muscular atrophy. *Muscle Nerve* 13:667-674, 1990.
- Luciano CA, Sivakumar S, Spector SA, et al: Reinnervation in clinically unaffected muscles of patients with prior paralytic poliomyelitis: Correlation between macro-electromyography and histology, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 394-396.
- Maselli RA, Cashman NR, Wolman EF, et al: Neuromuscular transmission as a function of motor unit size in patients with prior poliomyelitis. *Muscle Nerve* 15:648-655, 1992.
- Illa I, Leon-Monzon M, Dalakas MC: Regenerating and denervated human muscle fibers and satellite cells express N-CAM recognized by monoclonal antibodies to NK cells. *Ann Neurol* 31:46-52, 1992.
- Yunis EJ, Samaha FJ: Inclusion body myositis. *Lab Invest* 25:240-248, 1971.
- Pezeshkpour GH, Dalakas MC: Long-term changes in the spinal cord of patients with old poliomyelitis: Signs of continuous disease activity. *Arch Neurol* 45:505-508, 1988.
- Miller DC: Post-polio syndrome spinal cord pathology, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 186-193.
- Ginsberg AH, Gale MJ Jr, Rose LM, et al: T-cell alterations in late poliomyelitis. *Arch Neurol* 46:497-501, 1989.
- Johnson RT: Pathogenesis of poliovirus infections, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 361-365.
- Dalakas MC: Enteroviruses and human neuromuscular diseases, in Rotbard HA (ed): *Human Enterovirus Infections*. Washington, DC, ASM Press, 1995.
- Leon-Monzon M, Agboatwalla M, Dinsmore S, et al: Comparison of antibodies to the poliomyelitis virus in patients with acute paralytic poliomyelitis, post polio syndrome and amyotrophic lateral sclerosis. *Ann Neurol* 30:301-302, 1991.
- Sharief MK, Hentges R, Ciardi M: Intrathecal immune response in patients with the post-polio syndrome. *N Engl J Med* 325:749-755, 1991.
- Muir P, Nicholson F, Sharief MK, et al: Evidence for persistent enterovirus infection of the central nervous system in patients with previous paralytic poliomyelitis, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 219-232.
- Nelson KR: Creatine kinase and fibrillation potentials in patients with late sequelae of polio. *Muscle Nerve* 13:722-725, 1990.
- Leparc I, Kopecka H, Fuchs F, et al: Search for poliovirus in specimens from patients with the post-polio syndrome, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 233-236.
68. Allen GM, Gandevia SC, Neering IR, et al: Muscle performance, voluntary activation and perceived effort in normal subjects and patients with prior poliomyelitis. *Brain* 117:661-670, 1994.
69. Maselli RA, Wollmann R, Roos R: Function and ultrastructure of the neuromuscular junction in post polio syndrome, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 129-137.
70. Cashman NR, Trojan DA: Correlation of electrophysiology with pathology, pathogenesis and anticholinesterase therapy in post-polio, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 138-150.
71. Sharma KR, Kent-Braun J, Mynhier MA, et al: Excessive muscular fatigue in the postpoliomyelitis syndrome. *Neurology* 44:642-646, 1994.
72. Sivakumar K, Sinnwell T, Yildiz E, et al: Study of fatigue in muscles of patients with post-polio syndrome by *in vivo*  $^{31}\text{P}$  magnetic resonance spectroscopy: A metabolic cause for fatigue, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 397-401.
73. Parry GJ, Clarke S: Multifocal acquired demyelinating neuropathy masquerading as motor neuron disease. *Muscle Nerve* 11:103-107, 1988.
74. Parry GJ: Motor neuropathy with multifocal conduction block. *Semin Neurol* 13:269-275, 1993.
75. Sadiq SA, Thomas FP, Kilidireas K, et al: The spectrum of neurologic disease associated with anti-GM1 antibodies. *Neurology* 40:1067-1072, 1990.
76. Pestronk A, Chaudhury V, Feldman EL, et al: Lower motor neuron syndromes defined by patterns of weakness, nerve conduction abnormalities, and high titers of anti-glycolipid antibodies. *Ann Neurol* 27:316-326, 1990.
77. Dalakas MC: Polymyositis, dermatomyositis and inclusion body myositis. *N Engl J Med* 325:1487-1498, 1991.
78. Joy JL, Oh SJ, Baysal AI: Electrophysiological spectrum of inclusion body myositis. *Muscle Nerve* 13:949-951, 1990.
79. Lindberg C, Oldfors A, Hedstrom A: Inclusion body myositis: Peripheral nerve involvement. *J Neurol Sci* 99:327-338, 1990.
80. Spector SA, Gordon PL, Feuerstein IM, et al: Strength gains without muscle injury after strength training in patients with post polio muscular atrophy. *Muscle Nerve* 19:1282-1290, 1996.
81. Fillyaw MJ, Badger GJ, Goodwin GD, et al: The effects of long-term non-fatiguing resistance exercise in subjects with post-polio syndrome. *Orthopedics* 14:1253-1256, 1991.
82. Einarsson G: Muscle conditioning in late poliomyelitis. *Arch Phys Med Rehabil* 72:11-14, 1991.
83. Agre JC: The role of exercise in the patient with post-polio syndrome, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 321-334.
84. Bennett RL, Knowlton GC: Overwork weakness in partially denervated muscle. *Clin Orthop* 12:22-29, 1958.
85. Lenman JR: A clinical and experimental study of the effects of exercise on motor weakness in neurological disease. *J Neurol Neurosurg Psych* 22:182-194, 1959.
86. Treuth MS, Ryan AS, Pratley RE, et al: Effects of strength training on total and regional body composition in older men. *J Appl Physiol* 77:614-620, 1994.
87. Ryan AS, Treuth MS, Rubin MA, et al: Effects of strength training on bone mineral density: Hormonal and bone turnover relationships. *J Appl Physiol* 77:1678-1684, 1994.
88. Dalakas MC: How to design a therapeutic study in patients with the post-polio syndrome, in Dalakas MC, Bartfeld H, Kurland LT (eds): *The Post-Polio Syndrome: Advances in the Pathogenesis and Treatment*. New York, NY Academy of Sciences, 1995, pp 314-320.
89. Miller RG, Gelinas DF, Kent-Braun J, et al: The effect of recombinant insulin-like growth factor-1 upon exercise-induced fatigue and recovery in patients with post-polio syndrome. *Neurology* 48:A217, 1997.
90. Trojan DA, Cashman NR: An open trial of pyridostigmine in post-poliomyelitis syndrome. *Can J Neurol Sci* 22:223-227, 1995.
91. Jones DR, Speier J, Canine K, et al: Cardio-respiratory responses to aerobic training by patients with postpoliomyelitis sequelae. *JAMA* 261:3255-3258, 1989.
92. Owen RR, Jones D: Polio residuals clinic: Conditioning exercise program. *Orthopedics* 8:882-883, 1985.
93. Birk TJ: Poliomyelitis and the post-polio syndrome: Exercise capacities and adaptation: Current research, future directions, and widespread applicability. *Med Sci Sports Exerc* 25:466-472, 1993.
94. DiPietro L: Physical activity, body weight, and adiposity: An epidemiological perspective, in Holloszy JO (ed): *Exercise and Sport Sciences Reviews*. Baltimore, Williams and Wilkins, 1995, pp 275-304.
95. Waring WP, Maynard F, Grady W, et al: Influence of appropriate lower extremity orthotic management on ambulation, pain and fatigue in a postpolio population. *Arch Phys Med Rehabil* 70:371-375, 1989.
96. Perry J, Fleming C: Polio: Long-term problems. *Orthopedics* 8:877-881, 1985.
97. Forger NG, Roberts SL, Wong V, et al: Ciliary neurotrophic factor maintains motor neurons and their target muscles in developing rats. *J Neurosci* 13:4720-4726, 1993.
98. Sendtner M, Schmanbruch H, Stockli KA, et al: Ciliary neurotrophic factor prevents degeneration of motor neurons in mouse mutant progressive motor neuronopathy. *Nature* 358:502-504, 1992.
99. Lewis ME, Neff NT, Contreras PC, et al: Insulin-like growth factor-1: Potential for treatment of motor neuronal disorders. *Exp Neurol* 124:73-88, 1993.